

The Double-Lined Spectroscopic Binary Haro 1-14c

M. SIMON¹ and L.PRATO^{2,3}

ABSTRACT

We report detection of the low-mass secondary in the spectroscopic binary Haro 1-14c in the Ophiuchus star forming region. The secondary/primary mass ratio is 0.310 ± 0.014 . With an estimated photometric primary mass of $1.2 M_{\odot}$, the secondary mass is $\sim 0.4 M_{\odot}$ and the projected semi-major axis is ~ 1.5 AU. The system is well-suited for astrometric mapping of its orbit with the current generation of ground-based IR interferometers. This could yield precision values of the system's component masses and distance.

Subject headings: binaries: spectroscopic — stars: pre—main-sequence

1. Introduction

Reipurth et al. (2002) discovered that Haro 1-14c, a weak-lined pre—main-sequence star in the Ophiuchus star forming region (HBC 644, Herbig & Bell 1988), is actually the primary component of a single-lined spectroscopic binary (SB1) with a period of 591 days. We have been using high resolution infrared (IR) spectroscopy to detect the low-mass secondaries in main-sequence SB1s (Mazeh et al. 2003) and PMS SB1s (Prato et al. 2002a). We included Haro 1-14c in our program for two reasons. First, detection of its secondary would contribute to our study of the mass ratio distribution among PMS spectroscopic binaries (e.g., Prato et al. 2002a, 2002b). Second, it is important to test theoretical models of PMS evolution with stars whose masses have been measured independently (e.g., Simon et al. 2000; Hillenbrand & White 2004). Meaningful tests require masses known to a few percent; this is possible with masses measured dynamically. The fairly long period of Haro 1-14c and its location in the Ophiuchus SFR at ~ 140 pc distance suggested that the orbit might be resolvable with the current generation interferometers operating in the near IR. Mapping observations of Haro 1-14c, combined with the parameters derivable from analysis of the binary as a

¹Department of Physics and Astronomy, SUNY, Stony Brook, NY 11794-3800; michal.simon@sunysb.edu

²Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095-1562; lprato@astro.ucla.edu

³Current address: Lowell Observatory, 1400 West Mars Hill Road, Flagstaff, AZ 86001

double-lined spectroscopic binary (SB2), could yield precision measurements of the primary and secondary masses and the distance to the system (e.g., Boden et al. 2000). We have detected the secondary in Haro 1-14c in 6 observations distributed over its orbital period. This paper describes our results.

2. Observations and Data Reduction

We measured the radial velocities of the Haro 1-14c primary and secondary at the Keck II telescope of the Keck Observatory using the NIRSPEC high-resolution IR spectrometer (McLean et al. 1998, 2000). We centered the spectrometer at $1.55\ \mu\text{m}$ and obtained data following the procedures we have described previously (Prato et al. 2002a, 2002b). All the observations were made without the adaptive optics system and have average spectral resolution $R \sim 28,000$. The total integration times ranged from 6–20 minutes, depending on seeing conditions, which provided spectra with signal-to-noise ratio ~ 100 . The UT dates and modified Julian Dates (MJD) of the observations are listed in Table 1, columns 1 and 2. We used the REDSPEC software¹ to extract the spectra and place them on a wavelength scale. As in our previous work, we used only order 49 because it is almost completely free of terrestrial absorption lines.

To measure the radial velocities of the primary and secondary, we used the TODCOR two-dimensional algorithm (Zucker & Mazeh 1994) and our suite of spectral templates as we have described before (Mazeh et al. 2003, Prato et al. 2002a). Of the templates available, the main sequence star HR 8085, rotationally broadened to $12\ \text{km s}^{-1}$, usually provided the best fit to the primary of Haro 1-14c. Its K5 spectral type template is in reasonable agreement with the spectral type K3 listed in the Herbig and Bell Catalogue (Herbig & Bell 1988) for the component of Haro 1-14c brighter in visible light. The M1.5 main sequence star GL 15A, also rotationally broadened to $12\ \text{km s}^{-1}$, fit the secondary the best. The average secondary/primary flux ratio at $1.55\ \mu\text{m}$ was ~ 0.4 . However, because the main-sequence templates are not likely to match the surface gravity and metallicity of the PMS components of Haro 1-14c exactly, we do not regard as definitive the best-fitting spectral types and flux ratio. Table 1 lists our measured radial velocities of the primary and secondary, V_1 , and V_2 , and their estimated uncertainties.

Reipurth et al.’s (2002) observations of Haro 1-14c provided 56 measurements of its primary velocity and spanned ~ 3.6 orbital periods. These yielded accurate values of the binary center-of-mass velocity, γ , orbital period, P , velocity semi-amplitude of the primary

¹See <http://www2.keck.hawaii.edu/inst/nirspec/redspec/index.html>

K_1 , eccentricity, e , primary semi-major axis multiplied by the sine of the inclination, $a_1 \sin i$, mass function, $f(M)$, time of periastron passage, T_o , and longitude of the periastron of the primary, ω_1 . To these parameters, our measurements of the system contribute uniquely the velocity semi-amplitude of the secondary, K_2 , and hence the mass ratio, $q = M_2/M_1$, and secondary semi-major axis, $a_2 \sin i$. We merged our measurements with Reipurth et al's and, by least squares minimization of a phase shift of the SB2 data, first improved the precision of the P and T_o values enabled by the longer time span of the combined data. We then derived the other orbital parameters by least squares minimization; Table 2 lists the values and Figure 1 shows the corresponding double-lined orbital solution. The parameters in common with those calculated by Reipurth et al. are very close to their values, as expected, because the data set is dominated by their 56 measurements of the primary velocity. The improvement in precision is greatest for P and T_o because of the longer time interval of the merged data. We used the new values of P and T_o to calculate the phases of the SB2 observations listed in the last column of Table 1.

3. Discussion

The K3 spectral type of the primary (Reipurth et al. 2002) and PMS evolutionary tracks (e.g., Baraffe et al. 1998) for a few million year-old star suggest that $M_1 \sim 1.2 M_\odot$. This value, with the measured mass ratio, indicate that $M_2 \sim 0.4 M_\odot$. From Table 1 we have the estimate $a \sin i = (a_1 + a_2) \sin i = 229.0 \pm 8.0 \text{ Gm} \sim 1.5 \text{ AU}$. At the 140 pc assumed distance to the Ophiuchus SFR, the apparent angular size of the projected semi-major axis would be $\sim 11 \text{ mas}$.

Spectroscopic observations of SB2s leave three parameters undetermined: the two describing the binary orientation with respect to our line of sight, the orbital inclination, i , and the position angle of the line of nodes, and the distance to the binary. These parameters can be determined by resolving the components of the SB2 and mapping the orbit (e.g., Boden et al. 2000). The requirements for a binary to be resolvable with the current generation of IR interferometers are very stringent. The apparent angular extent of the binary orbit must be small enough that it lies within the diffraction limit of a single telescope of the interferometer, but large enough that it is resolvable at the projected interferometer baselines. Both components of the binary must be bright enough in the near-IR, usually K-band, that fringes can be detected, and bright enough in the visible for wavefront correction by adaptive optics. Targets satisfying these requirements are rare.

With $K \sim 7.8 \text{ mag}$ (2MASS catalog) and $V \sim 12.3 \text{ mag}$ (Rydgren et al. 1976), and an estimated projected semi-major axis of 11 mas, Haro 1-14c is suitable study for study by the

combined techniques of IR interferometry and spectroscopy. The measurements will yield the component masses, system distance (hence total luminosity), component flux ratios at the wavelengths of the interferometric and spectroscopic observations (enabling apportionment of the luminosity between the components), and estimates of the effective temperature (from the best matching templates for the analysis of the spectra). These parameters, together with the reasonable assumption of equal age of the components, will enable definitive tests of theoretical calculations of PMS evolution at masses below $0.5 M_{\odot}$ (e.g. Simon et al. 2000; Hillenbrand and White 2004).

4. Conclusions

We have detected the secondary in the PMS binary Haro 1-14c by high spectral resolution observations at $1.55\mu\text{m}$. We find that:

1) The mass ratio of ~ 0.31 suggests that for an estimated photometric mass of $1.2 M_{\odot}$ for the primary, the secondary mass is $\sim 0.4 M_{\odot}$.

2) Assuming these estimated masses, the projected orbital semi-major axis is $a \sin i \sim 1.5$ AU. At an assumed 140 pc distance to the Ophiuchus SFR, the binary is angularly resolvable with the current generation of interferometers operating in the near-IR.

3) Astrometric measurements that map the binary orbit, and continued spectroscopic observations that improve the precision of the mass ratio offer the promise to measure the distance to the binary and the masses of its components at the few percent level.

5. Acknowledgements

We thank the referee for a constructive comment that helped us improve the discussion section. We are very grateful to Tsevi Mazeh and Shay Zucker for allowing us to use their TODCOR algorithm. We thank the Keck observing assistants, and the staff in Waimea, for their thorough support. This research was supported in part by NSF Grant AST 02-05427. The data were obtained at the W.M. Keck Observatory which is operated as a scientific partnership between the California Institute of Technology, the University of California, and NASA. The observatory was made possible by the generous financial support of the W.M. Keck Foundation. Our research made use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. Our research

also used the SIMBAD database, operated at CDS, Strasbourg, France. The authors extend special thanks to those of Hawaiian ancestry on whose sacred mountain we are privileged to be guests.

REFERENCES

- Baraffe, I., Chabrier, G., Allard, F., and Hauschildt, P. H. 1998, *A&A*, 337, 403
- Boden, A.F., Creech-Eakman, M.J., & Queloz, D. 2000, *ApJ*, 536, 880
- Herbig, G.H. & Bell, K.R. 1988, *Lick Obs. Bull.* No. 1111
- Hillenbrand, L.A. & White, R.J., 2004, *ApJ*, 604, 741
- Mazeh, T., Simon, M., Prato, L., Markus B., & Zucker, S. 2003, *ApJ*, 599, 1344
- McLean, I. S., et al. 1998, *SPIE*, 3354, 566
- McLean, I. S., et al. 2000, *SPIE*, 4008, 1048
- Prato, L., Simon, M., Mazeh, T., McLean, I. S., Norman, & D., Zucker, S. 2002a, *ApJ*, 569, 863
- Prato, L., Simon, M., Mazeh, T., S. Zucker, & McLean, I. S. 2002b, *ApJ*, 579, L99
- Reipurth, B., Lindgren, H., Mayor, M., Mermilliod, J-C., & Cramer, N. 2002, *AJ*, 124, 2813
- Rydgren, A.E., Strom, S.E., & Strom, K.M. 1976, *ApJS*, 30, 307
- Simon, M., Dutrey, A., & Guilloteau, S. 2000, *ApJ*, 545, 1034
- Wilson, O.C. 1941, *ApJ*, 93, 29
- Zucker, S., & Mazeh, T. 1994, *ApJ*, 420, 806




Fig. 1.— Radial velocity as a function of phase for Haro 1-14c. The crosses represent Reipurth et al.’s (2002) measurements of the primary velocity. The open symbols, and 1σ uncertainties indicate our measurements for the primary (circles) and secondary (squares). The solid line shows the best fit to the primary star data, and the dashed line for the secondary. The horizontal dashed line indicates the center-of-mass velocity of the system. The abscissa spans a 20% redundancy in phase for clarity.

Table 1. Measured Velocities

UT Date	MJD	V_1 km s ⁻¹	σ km s ⁻¹	V_2 km s ⁻¹	σ_2 km s ⁻¹	Phase
2001 June 1	52061.50	-2.9	2.0	-18.7	2.0	0.307
2002 July 17	52472.28	-16.1	2.0	18.7	2.0	0.002
2003 Feb 8	52678.67	-4.9	2.0	-18.3	2.0	0.351
2003 Aug 10	52861.32	-8.4	2.0	-5.8	2.0	0.660
2003 Sep 8	52890.20	-9.3	2.0	-3.6	2.0	0.709
2004 Jan 26	53030.69	-19.4	2.0	26.6	2.0	0.947

Table 2. Orbital Elements

Parameter	Value
γ	-8.71 ± 0.07 km s ⁻¹
P	591.3 ± 0.3 days
K_1	8.52 ± 0.13 km s ⁻¹
K_2	27.5 ± 1.3 km s ⁻¹
e	0.617 ± 0.008
q	0.310 ± 0.014
$a_1 \sin i$	54.2 ± 0.8 Gm
$a_2 \sin i$	174.8 ± 7.8 Gm
$M_1 \sin^3 i$	0.98 ± 0.14 $M_\odot$
$M_2 \sin^3 i$	0.30 ± 0.05 $M_\odot$
ω_1	$232.90 \pm 0.55^\circ$
T_0	45375.4 ± 3.0 MJD